

Hypergeometric Integrals of Motion and Affine Gaudin Models

Charles Young


University of Hertfordshire, U.K.

SEMPS, Surrey,
March 2018

based on work with **Sylvain Lacroix** and **Benoît Vicedo**: [1804.01480 and to appear]

Quantum Gaudin Model

- ▶ Let \mathfrak{g} be any symmetrizable Kac-Moody algebra (e.g. \mathfrak{sp}_M , or $\widehat{\mathfrak{sl}}_M, \dots$)
- ▶ Assign irreducible highest-weight \mathfrak{g} -modules $\{L_{\lambda_i}\}$ to marked points $\{z_i\}$ in \mathbb{C} :



- ▶ Canonical element: $\Xi = \sum_{\text{roots } \alpha} \Xi_{\alpha}, \quad \Xi_{\alpha} \in \mathfrak{g}_{\alpha} \otimes \mathfrak{g}_{-\alpha}$

$$\mathcal{H}_i = \sum_{\substack{j=1 \\ j \neq i}}^N \frac{\Xi^{(ij)}}{z_i - z_j} \in U(\mathfrak{g})^{\otimes N}, \quad i = 1, \dots, N$$

Quadratic Gaudin Hamiltonians $\mathcal{H}_i : \bigotimes_{k=1}^N L_{\lambda_k} \rightarrow \bigotimes_{k=1}^N L_{\lambda_k}$

Bethe ansatz for Gaudin models

- ▶ Gaudin model solvable by a form of Bethe ansatz:
 - ▶ Pick $m \geq 0$ additional marked points w_j (Bethe roots)
 - ▶ Associate to each a simple root $\alpha_{c(j)}$
 - ▶ Construct Bethe vector $\psi = \psi(\{z_i\}, \{\lambda_i\}, \{w_j\}, \{\alpha_{c(j)}\})$
- ▶ **Theorem:** if Bethe roots $\{w_j\}$ obey **Bethe equations** then ψ is a joint eigenvector of the \mathcal{H}_i , with **explicit eigenvalues**.

$$-\sum_{i=1}^N \frac{(\lambda_i | \alpha_{c(j)})}{w_j - z_i} + \sum_{\substack{i=1 \\ i \neq j}}^m \frac{(\alpha_{c(i)} | \alpha_{c(j)})}{w_j - w_i} = 0, \quad j = 1, \dots, m.$$
$$E_i := \sum_{\substack{j=1 \\ j \neq i}}^N \frac{(\lambda_i | \lambda_j)}{z_i - z_j} - \sum_{j=1}^m \frac{(\lambda_i | \alpha_{c(j)})}{z_i - w_j}, \quad i = 1, \dots, N.$$

- ▶ Theorem holds for **any** symmetrizable Kac-Moody algebra \mathfrak{g} .
(General proof is in terms of hyperplane arrangements [Schechtman & Varchenko, 91])

Finite type \mathfrak{g} – Gaudin algebra and Opers

For \mathfrak{g} of finite type (i.e. one of $\mathfrak{sl}_M, \mathfrak{so}_M, \mathfrak{sp}_M, \mathfrak{e}_6, \mathfrak{e}_7, \mathfrak{e}_8, \mathfrak{f}_4, \mathfrak{g}_2$)
much more is known:

$$\mathcal{H}_i \in \mathcal{B} \subset U(\mathfrak{g})^{\otimes N}$$

Gaudin/Bethe Algebra

[Feigin Frenkel Reshetikhin]
[Mukhin Tarasov Varchenko]

- ▶ \mathcal{B} : a (large) commutative subalgebra of $U(\mathfrak{g})^{\otimes N}$
generated by \mathcal{H}_i together with **higher Gaudin Hamiltonians**
- ▶ ψ is a joint eigenvector for the entire algebra \mathcal{B}
- ▶ **Theorem:** Joint eigenvalues encoded as functions on a space of **opers**
[Frenkel], [Rybnikov], [Mukhin Tarasov Varchenko]
– “Geometric Langlands correspondence”

Main questions:

Suppose \mathfrak{g} is of untwisted **affine type** (e.g. $\widehat{\mathfrak{sl}}_M, \widehat{\mathfrak{so}}_M, \widehat{\mathfrak{sp}}_M, \dots$)

1. Are there higher Gaudin Hamiltonians?
2. If yes, then what parameterizes their eigenvalues?
Functions onopers?? Whatopers?? What do such functions look like??

...important questions for mathematical physics because affine (quantum) Gaudin models are closely related to integrable (quantum) field theories in $1 + 1$ dimensions

Plan of this talk:

- (i) Define a notion of affineopers, generalizing definitions from finite type in the most direct way possible.
- (ii) Main result: the functions on the space of affineopers are of a very different character than in the finite case:
they are given by **hypergeometric-type integrals** over cycles of a twisted homology defined by the levels of the modules at the marked points.
- (iii) Conjecture: these integrals are the eigenvalues of (higher) Gaudin Hamiltonians
(... prompts a conjecture about the form of such Hamiltonians themselves)
- (iv) Check this conjecture in some special cases (use GKO coset construction/Integrals of Motion of quantum KdV and quantum Boussinesq theory)

Review: Operers and Miura operers in finite types

Suppose \mathfrak{g} is of finite type. Let ${}^L\mathfrak{g}$ be its Langlands dual (also of finite type).

- ▶ Cartan decomposition: ${}^L\mathfrak{g} = {}^L\mathfrak{n}_- \oplus {}^L\mathfrak{h} \oplus {}^L\mathfrak{n}_+$
- ▶ Chevalley generators: $\check{f}_i, \check{e}_i, i = 1, \dots, \ell$.
- ▶ Simple coroots: $\alpha_i := [\check{e}_i, \check{f}_i]$ (are the simple roots of \mathfrak{g})

Definition: A Miura ${}^L\mathfrak{g}$ -oper is a connection of the form

$$\nabla = d + \left(\bar{p}_{-1} + u(z) \right) dz.$$

Principal nilpotent element

$$\bar{p}_{-1} := \sum_{i=1}^{\ell} \check{f}_i$$

rational function valued in Cartan ${}^L\mathfrak{h} \cong \mathfrak{h}^*$

For us, $u(z)$ is of the form

$$u(z) = - \sum_{i=1}^N \frac{\lambda_i}{z - z_i} + \sum_{j=1}^m \frac{\alpha_{c(j)}}{z - t_j}$$

and encodes the marked points $\{z_i\}$, Bethe roots $\{w_j\}$, highest weights $\{\lambda_i\}$ and “colours” of the Bethe roots $\{c(j)\}$.

Definition: An $L\mathfrak{g}$ -oper is a **gauge equivalence class** $[\nabla]$ of connections of the form

$$\nabla = d + \left(\bar{p}_{-1} + b(z) \right) dz$$

rational function valued in Borel $L\mathfrak{b}_+ \cong L\mathfrak{h} \oplus L\mathfrak{n}_+$

under the gauge action of the unipotent subgroup ${}^L N = \exp({}^L \mathfrak{n}_+)$.

Fact: Each oper $[\nabla]$ has a unique representative of **canonical form**

$$d + \left(\bar{p}_{-1} + \sum_{r \in \bar{E}} \bar{v}_r(z) \bar{p}_r \right) dz.$$

finite (multi)set of exponents of $L\mathfrak{g}$

rational coefficient functions

element $\bar{p}_r \in L\mathfrak{n}_+$ of grade r in principal gradation (h.w. vector for principal \mathfrak{sl}_2)

Corollary: These $\bar{v}_r(z)$ are “good coordinates” on the space ofopers.

- ▶ Each Miura oper ∇ defines an underlying oper $[\nabla]$

Fact: The Bethe equations precisely ensure $\bar{v}_r(z)$ have poles only at the marked points z_1, \dots, z_N (and ∞) and not at the Bethe roots w_1, \dots, w_m .

Dictionary:

Miura oper $\nabla \longleftrightarrow u(z) \in {}^L\mathfrak{h} \longleftrightarrow$ joint eigenvector ψ of Gaudin Hamiltonians

Underlying oper $[\nabla] \longleftrightarrow \{\bar{v}_r(z) \in \mathbb{C}\}_{r \in \bar{E}} \longleftrightarrow$ eigenvalues of all Gaudin Hamiltonians

Opers and Miura opers in affine types

Suppose \mathfrak{g} is of **untwisted affine** type. Let ${}^L\mathfrak{g}$ be Langlands dual (**affine**, maybe twisted).

- ▶ Cartan decomposition: ${}^L\mathfrak{g} = {}^L\mathfrak{n}_- \oplus {}^L\mathfrak{h} \oplus {}^L\mathfrak{n}_+$
- ▶ Chevalley generators: $\check{f}_i, \check{e}_i, i = 0, 1, \dots, \ell$; coroots $\alpha_i = [\check{e}_i, \check{f}_i]$

Definition: A **Miura ${}^L\mathfrak{g}$ -oper** is a connection of the form

$$\nabla = d + \left(p_{-1} + u(z) \right) dz.$$

Principal **nilpotent** element

$$p_{-1} := \sum_{i=0}^{\ell} \check{f}_i$$

rational function valued in Cartan ${}^L\mathfrak{h} \cong \mathfrak{h}^*$

- ▶ $u(z)$ as before – except ‘colours’ of Bethe roots $c(j) \in \{0, 1, \dots, \ell\}$ can include 0.
- ▶ **Principal derivation element:** $\rho \in {}^L\mathfrak{h}$. $[\rho, \check{e}_i] = \check{e}_i$, $[\rho, \check{f}_i] = -\check{f}_i$.
- ▶ Decompose $u(z)$ in basis $\{\alpha_i\}_{i=0}^{\ell} \cup \{\rho\}$:

$$\nabla = d + \left(p_{-1} - \frac{\varphi(z)}{h^\vee} \rho + \sum_{i=0}^{\ell} u_i(z) \alpha_i \right) dz, \quad \varphi(z) = \sum_{i=1}^N \frac{k_i}{z - z_i}$$

where $k_i = \langle k, \lambda_i \rangle$ are the levels of the L_{λ_i} . Call $\varphi(z)$ the **twist function**.

Definition: An $L\mathfrak{g}$ -oper is a **gauge equivalence class** $[\nabla]$ of connections of the form

$$\nabla = d + \left(p_{-1} + b(z) \right) dz$$

rational function valued in Borel $L\mathfrak{b}_+ \cong L\mathfrak{h} \oplus L\mathfrak{n}_+$

under the gauge action of the **unipotent** subgroup ${}^L N = \exp({}^L \mathfrak{n}_+)$.

Theorem: [Lacroix, Vicedo, CY]

(i) Each oper $[\nabla]$ has a **unique** representative of **quasi-canonical form**

$$d + \left(p_{-1} - \frac{\varphi(z)}{h^\vee} \rho + \sum_{r \in E} v_r(z) p_r \right) dz.$$

countably infinite (multi)set of exponents of $L\mathfrak{g}$

rational coefficient functions

element $p_r \in L\mathfrak{n}_+$ of
grade r in principal gradation
(\in principal Heisenberg subalgebra)

(ii) The functions $\varphi(z)$ and $v_1(z)$ are unique. But the functions $v_r(z)$, $r \geq 2$, are unique only up to transformations of the form

$$v_r(z) \mapsto v_r(z) - g'_r(z) + \frac{r\varphi(z)}{h^\vee} g_r(z)$$

for any rational functions $g_r(z)$.

Corollary: ~~These $v_r(z)$ are “good coordinates” on the space of affine ops.~~

... so how to construct well-defined functions on the space of affine ops?

- ▶ Define multivalued function $\mathcal{P}(z) := \prod_{i=1}^N (z - z_i)^{k_i}$ whose log-derivative is $\varphi(z)$.
- ▶ Gauge freedom in $v_r(z)$ is then


$$\mathcal{P}(z)^{-r/h^\vee} v_r(z) \mapsto \mathcal{P}(z)^{-r/h^\vee} v_r(z) - \partial_z (\mathcal{P}(z)^{-r/h^\vee} g_r(z)).$$

- ▶ To get gauge-invariant quantities we should consider **integrals** of $\mathcal{P}(z)^{-r/h^\vee} v_r(z)$...

integrals over any cycle γ which is not only closed but also around which \mathcal{P} is single-valued. . .

$$I_\gamma^{(r)} := \int_\gamma \mathcal{P}(z)^{-r/h^\vee} v_r(z) dz$$

Prototypical example of such cycles are Pochhammer contours



Corollary: These integrals $I_\gamma^{(r)}$ are well-defined functions on the space of affineopers.

Proposition: The Bethe equations precisely ensure **there exists a gauge in which** $\{v_r(z)\}_{r \in E}$ have poles only at the marked points z_1, \dots, z_N (and ∞) and not at the Bethe roots w_1, \dots, w_m .

Conjectures

1. These integrals $I_\gamma^{(r)}$ are the eigenvalues of higher affine Gaudin Hamiltonians.
2. The Hamiltonians themselves are integrals,

$$H_\gamma^{(r)} := \int_\gamma \mathcal{P}(z)^{-r/h^\vee} S_r(z)_0 dz$$

for certain “densities” $S_r(z)_0 \in \hat{U}(\mathfrak{g}^{\oplus N})$ depending rationally on z .

In particular, each Hamiltonian is labelled by

- ▶ an exponent r from the infinite multiset E of exponents and
- ▶ a contour γ (more precisely, an element of a twisted homology space)

Checks

- ▶ Semiclassics
- ▶ Cubic Hamiltonians
- ▶ GKO coset constructions (2-point Gaudin models for $\widehat{\mathfrak{sl}}_2$ and $\widehat{\mathfrak{sl}}_3$)

Semiclassics

Recall results on classical Principal Chiral Models (PCMs)

[Evans, Hassan, MacKay, Mountain]

- ▶ Let $j_+ = g^{-1}\partial_+g$ where $g = g(x, t) \in G$ is the PCM field.
- ▶ There are Poisson-commuting conserved charges of the form

$$\int_0^{2\pi} dx K_{ab\dots c} j_+^a j_+^b \dots j_+^c.$$

Here $K_{ab\dots c}$ are certain invariant tensors whose degrees $\in \{ \text{exponents of } G \text{ repeating modulo the Coxeter number} \} = \{ \text{the exponents of the affine algebra} \}$

- ▶ Classical PCMs can be interpreted as **classical** affine Gaudin models and then these conserved charges are of the form

[Vicedo],[Lacroix, Magro, Vicedo]

$$\int_0^{2\pi} dx K_{ab\dots c} L(z_{(0)})^a L(z_{(0)})^b \dots L(z_{(0)})^c.$$


where $L(z)$ is the (Gaudin) Lax matrix and $z_{(0)}$ is a **zero of the twist function** $\varphi(z)$.

Semiclassics

On the other hand, one can re-introduce \hbar in the quantum-mechanical constructions above:

$$H_\gamma^{(r)} = \int_\gamma \mathcal{P}(z)^{-r/(\hbar h^\vee)} S_r^{(\hbar)}(z) dz$$

Then in the $\hbar \rightarrow 0$ limit, deform contour γ to apply method of steepest descents:



Integrals of the form $H_\gamma^{(r)}$ localize at the saddle points of $\mathcal{P}(z) =$ **zeros of $\varphi(z)$!**

(And count of zeros (= $N - 1$) agrees with count of independent cycles.)

(Reminiscent of passage from KZ equations to Gaudin model – yet conceptually quite separate)


Cubic Hamiltonians

- ▶ Simplest general direct check is in types $\widehat{\mathfrak{sl}}_M$ with $M \geq 3$.
- ▶ Check for $r = 1, 2$ only so far, i.e. quadratic and cubic Hamiltonians.
- ▶ (Guess that) densities $S_r(z)_0$ are actually Fourier **zero modes** of certain **states** in tensor product of Vacuum verma modules $\mathbb{V}_0^k = \bigotimes_{i=1}^N \mathbb{V}_0^{k_i}$

$$S_1(z) := \frac{1}{2} I_{-1}^a(z) I_{-1}^a(z) |0\rangle^k,$$

$$S_2(z) := \frac{1}{3} t_{abc} I_{-1}^a(z) I_{-1}^b(z) I_{-1}^c(z) |0\rangle^k,$$

where $I_{-1}^a(z) := \sum_{i=1}^N \frac{I_{-1}^{a, (\text{tensor factor } i)}}{z - z_i}$ and $|0\rangle^k = |0\rangle \otimes \dots \otimes |0\rangle$.



Theorem: [Lacroix,Vicedo,CY] For $i, j \in \{1, 2\}$,

$$S_i(z)_{(0)}S_j(w) = D_z^{(i)}A_{ij}(z, w) + D_w^{(j)}B_{ij}(z, w) + TC_{ij}(z, w),$$

for some \mathbb{V}_0^k -valued rational functions $A_{ij}(z, w)$, $B_{ij}(z, w)$ and $C_{ij}(z, w)$.

Proof Direct (lengthy) calculation... e.g.

$$\begin{aligned} A_{22}(z, w) &= \left(\frac{2h^{\vee 3} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^3} I_{-4}^a(z) I_{-1}^a(z) - \frac{4h^{\vee 3} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^3} I_{-4}^a(z) I_{-1}^a(w) \right. \\ &\quad - \frac{2h^{\vee 2} \left(1 - \frac{4}{h^{\vee 2}}\right) \varphi(z)}{(z-w)^2} I_{-4}^a(z) I_{-1}^a(w) - \frac{h^{\vee 3} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^2} I_{-4}^a(z) I_{-1}^a(w) \\ &\quad - \frac{2h^{\vee 3} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^3} I_{-3}^a(z) I_{-2}^a(z) + \frac{2h^{\vee 2} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^2} f_{abc} I_{-3}^a(z) I_{-1}^b(z) I_{-1}^c(w) \left. C_{22}(z, w) \right. \\ &\quad \left. + \frac{h^\vee}{z-w} t_{abc} t_{cde} I_{-2}^a(z) I_{-1}^b(z) I_{-1}^c(w) I_{-1}^d(w) \right) |0\rangle^k \\ B_{22}(z, w) &= \left(- \frac{2h^{\vee 3} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^3} I_{-4}^a(z) I_{-1}^a(z) + \frac{8h^{\vee 3} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^3} I_{-4}^a(z) I_{-1}^a(w) \right. \\ &\quad + \frac{4h^{\vee 2} \left(1 - \frac{4}{h^{\vee 2}}\right) \varphi(z)}{(z-w)^2} I_{-4}^a(z) I_{-1}^a(w) - \frac{2h^{\vee 2} \left(1 - \frac{4}{h^{\vee 2}}\right) \varphi(w)}{(z-w)^2} I_{-4}^a(z) I_{-1}^a(w) \\ &\quad + \frac{h^{\vee 3} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^2} I_{-4}^a(z) I_{-1}^a(w) + \frac{2h^{\vee 3} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^3} I_{-3}^a(z) I_{-2}^a(z) \\ &\quad - \frac{2h^{\vee 2} \left(1 - \frac{4}{h^{\vee 2}}\right)}{(z-w)^2} f_{abc} I_{-3}^a(z) I_{-1}^b(z) I_{-1}^c(w) \\ &\quad \left. - \frac{h^\vee}{z-w} t_{abc} t_{cde} I_{-2}^a(z) I_{-1}^b(z) I_{-1}^c(w) I_{-1}^d(w) \right) |0\rangle^k \end{aligned}$$

Corollary: The corresponding Hamiltonians, i.e. contour integrals of zero modes, commute.

GKO coset construction and qKdV integrals of motion

Consider Gaudin model for $\widehat{\mathfrak{sl}}_2$ with 2 marked points.

Quadratic Hamiltonian:

$$\mathcal{H} := \mathcal{H}_1 = -\mathcal{H}_2 = \frac{\Xi}{z_1 - z_2} \quad \text{where} \quad \Xi = d \otimes k + k \otimes d + \sum_n I_n^a \otimes I_{a,-n}$$

On the other hand, have Segal-Sugawara generators of Virasoro algebra at sites 1 and 2, and the diagonal copy:

$$T^{(1)}(x) := \frac{1}{2(k_1 + h^\vee)} \sum_{n \in \mathbb{Z}} : I_n^{a(1)} I_{a,-n}^{(1)} : \quad T^{(2)}(x) := \frac{1}{2(k_2 + h^\vee)} \sum_{n \in \mathbb{Z}} : I_n^{a(2)} I_{a,-n}^{(2)} :$$

$$T^{(diag)}(x) := \frac{1}{2(k_1 + k_2 + h^\vee)} \sum_{n \in \mathbb{Z}} : (I_n^{a(1)} + I_n^{a(2)}) (I_{a,-n}^{(1)} + I_{a,-n}^{(2)}) :$$

And then the Goddard-Kent-Olive coset generators of Virasoro are:

$$T^{(GKO)}(x) := T^{(1)}(x) + T^{(2)}(x) - T^{(diag)}(x) =: \sum_{n \in \mathbb{Z}} L_n x^{-n-2}$$

Fact: The quadratic Gaudin Hamiltonian is the GKO Virasoro zero mode:

$$\Xi = -(k_1 + k_2 + h^\vee) L_0$$

But the Virasoro algebra is known to have a large commutative subalgebra, called the algebra of **Quantum Integrals of Motion** (QIMs) (of quantum (m)KdV).

[Sasaki, Yamanaka],[Feigin, Frenkel]

$$I_1 = L_0$$

$$I_3 = 2 \sum_{n=1}^{\infty} L_{-n} L_n + L_0^2 - \frac{c+2}{12} L_0 + \frac{c(5c+22)}{2880}$$

$$I_5 = \dots$$

Since the first of these is the quadratic Gaudin Hamiltonian, have natural:

Conjecture/Definition: [Feigin, Frenkel] In this case (2 sites, $\widehat{\mathfrak{sl}}_2$) the higher Quantum Integrals of Motion are the higher affine Gaudin Hamiltonians.

Taking this as a definition, have an arena to test conjecture about eigenvalues. . .

\mathbb{C}

$z_1 = 0$ $z_2 = 1$

Top multiplicity space = Virasoro module with

$$c(a, b) = 1 - \frac{6}{(a+b+2)(a+b+3)}$$

$$\Delta(a, b) = \frac{b(b+2)}{4(a+b+2)(a+b+3)}$$

$$L_{a\Lambda_0 + b\Lambda_1} \otimes L_{\Lambda_0} = L_{(a+1)\Lambda_0 + b\Lambda_1} \otimes \mathcal{U} \oplus \dots$$

- ▶ Virasoro calculation: Vacuum value of, e.g. I_5 is

$$I_5 = \Delta^3 - \frac{c+4}{8} \Delta^2 + \frac{(c+2)(3c+20)}{576} \Delta + \frac{(-c)(3c+14)(7c+68)}{290304}$$

- ▶ Affine oper calculation: $u(z) := \frac{\frac{1}{4}(b-a)}{z} - \frac{\frac{1}{4}}{z-1}$, $\varphi(z) := \frac{a+b}{z} + \frac{1}{z-1}$ and

$$I_\gamma^{(5)} = \int_\gamma \mathcal{P}(z)^{-5/2} v_5(z) dz$$

where $v_5(z)$ is given by

$$\begin{aligned} & \frac{u(z)^2 \left(\frac{d^3}{dz^3} \varphi(z)\right)}{16} + \frac{5u(z) \left(\frac{d}{dz} u(z)\right) \left(\frac{d^2}{dz^2} \varphi(z)\right)}{16} + \frac{-11u(z)^2 \varphi(z) \left(\frac{d^2}{dz^2} \varphi(z)\right)}{16} + \frac{-7u(z)^2 \left(\frac{d}{dz} \varphi(z)\right)^2}{16} + \frac{5u(z) \left(\frac{d^2}{dz^2} u(z)\right) \left(\frac{d}{dz} \varphi(z)\right)}{8} \\ & + \frac{-45u(z) \varphi(z) \left(\frac{d}{dz} u(z)\right) \left(\frac{d}{dz} \varphi(z)\right)}{16} + \frac{23u(z)^2 \varphi(z)^2 \left(\frac{d}{dz} \varphi(z)\right)}{8} + \frac{-7u(z)^4 \left(\frac{d}{dz} \varphi(z)\right)}{16} + \frac{-u(z) \left(\frac{d^4}{dz^4} u(z)\right)}{16} + \frac{5u(z) \varphi(z) \left(\frac{d^3}{dz^3} u(z)\right)}{8} + \frac{-35u(z) \varphi(z)^2 \left(\frac{d}{dz} \varphi(z)\right)}{16} \\ & + \frac{7u(z)^3 \left(\frac{d^2}{dz^2} u(z)\right)}{16} + \frac{11u(z)^2 \left(\frac{d}{dz} u(z)\right)^2}{16} + \frac{25u(z) \varphi(z)^3 \left(\frac{d}{dz} u(z)\right)}{8} + \frac{-43u(z)^3 \varphi(z) \left(\frac{d}{dz} u(z)\right)}{16} + \frac{-3u(z)^2 \varphi(z)^4}{2} + \frac{25u(z)^4 \varphi(z)^2}{16} + \frac{-u(z)}{8} \end{aligned}$$

- ▶ Result: up to constants independent of the \mathfrak{sl}_2 weight b , both I_5 and $I_\gamma^{(5)}$ are equal to

$$425k^6 + 6375k^5 - 2898b^2k^4 - 5796bk^4 + 36287k^4 - 28980b^2k^3 - 57960bk^3 + 97245k^3 + 3780b^4k^2 + 15120b^3k^2 - 84042b^2k^2 - 198324bk^2 + 121724k^2 + 18900b^4k + 75600b^3k - 57960b^2k - 267120bk + 57120k - 1512b^6 - 9072b^5 + 60480b^3 + 12096b^2 - 120960b$$

- ▶ Similar checks works with (up to 2) Bethe roots instead of vacuum.
- ▶ Also have Cubic Affine Gaudin Hamiltonian, so can also try $\widehat{\mathfrak{sl}}_3$ case:

$\widehat{\mathfrak{sl}}_3$ -Coset construction of W_3 algebra

\mathbb{C}

$$z_1 = 0 \quad z_2 = 1$$

Top multiplicity space = W_3 module
[Bais, Bouwknegt, Surridge, Schoutens]

$$L_{a\Lambda_0 + b\Lambda_1 + c\Lambda_2} \otimes L_{\Lambda_0} = L_{(a+1)\Lambda_0 + b\Lambda_1 + c\Lambda_2} \otimes \mathcal{U} \oplus \dots$$

- ▶ On specializing to case of 2 points and $\widehat{\mathfrak{sl}}_M$, find $T = \int_{\gamma} \mathcal{P}(z)^{-1/3} S_1(z) dz$ and

$$\begin{aligned} W &= \int_{\gamma} \mathcal{P}(z)^{-2/3} S_2(z) dz \\ &\propto \frac{1}{3} t_{abc} I_{-1}^{a(1)} I_{-1}^{b(1)} I_{-1}^{c(1)} |0\rangle^{\mathbf{k}} \left(-\frac{2}{M} k_2\right) \left(-\frac{2}{M} k_2 - 1\right) \left(-\frac{2}{M} k_2 - 2\right) \\ &\quad + t_{abc} I_{-1}^{a(1)} I_{-1}^{b(1)} I_{-1}^{c(2)} |0\rangle^{\mathbf{k}} \left(-\frac{2}{M} k_1 - 2\right) \left(-\frac{2}{M} k_2 - 1\right) \left(-\frac{2}{M} k_2 - 2\right) \\ &\quad + t_{abc} I_{-1}^{a(1)} I_{-1}^{b(2)} I_{-1}^{c(2)} |0\rangle^{\mathbf{k}} \left(-\frac{2}{M} k_1 - 1\right) \left(-\frac{2}{M} k_1 - 2\right) \left(-\frac{2}{M} k_2 - 2\right) \\ &\quad + \frac{1}{3} t_{abc} I_{-1}^{a(2)} I_{-1}^{b(2)} I_{-1}^{c(2)} |0\rangle^{\mathbf{k}} \left(-\frac{2}{M} k_1\right) \left(-\frac{2}{M} k_1 - 1\right) \left(-\frac{2}{M} k_1 - 2\right) \end{aligned}$$

are the coset conformal and W vectors.

- ▶ Quantum Integrals of Motion $I_1, I_2, I_4, I_5, I_7, I_8, \dots$ are known.
- ▶ Check at least vacuum values of I_2, I_4, I_5 .

Open questions

- ▶ Existence proof (or explicit formula) for the higher Hamiltonians?
- ▶ Relation to ODE/IM. [Bazhanov, Lukyanov, Zamilodchikov] [Dorey, Dunning, Tateo] ? (“dual”?)
- ▶ Relation to Integrals of Motion in quantum toroidal algebras? [Feigin, Jimbo, Mukhin]