

APPROXIMATING THE MAXIMUM ERGODIC AVERAGE VIA PERIODIC ORBITS

D. COLLIER AND I.D. MORRIS

ABSTRACT. Let $\sigma: \Sigma_A \rightarrow \Sigma_A$ be a subshift of finite type, let \mathcal{M}_σ be the set of all σ -invariant Borel probability measures on Σ_A , and let $f: \Sigma_A \rightarrow \mathbb{R}$ be a Hölder continuous observable. There exists at least one σ -invariant measure μ which maximises $\int f d\mu$. The following question was asked by B.R. Hunt, E. Ott and G. Yuan: how quickly can the maximum of the integrals $\int f d\mu$ be approximated by averages along periodic orbits of period less than p ? We give an example of a Hölder observable f for which this rate of approximation is slower than stretched-exponential in p .

AMS subject classification: 37B10, 37D20, 49J30.

1. INTRODUCTION

Let $\sigma: \Sigma_A \rightarrow \Sigma_A$ be a subshift of finite type, let \mathcal{M}_σ be the set of all σ -invariant Borel probability measures on Σ_A , and let $f: \Sigma_A \rightarrow \mathbb{R}$ be Hölder continuous. There is at least one invariant measure $\mu \in \mathcal{M}_\sigma$, which we term a *maximising measure* for f , such that

$$\int f d\mu = \beta(f) := \sup_{\nu \in \mathcal{M}_\sigma} \int f d\nu.$$

In this paper we investigate the problem of finding invariant measures supported on periodic orbits which approximately realise this maximum. More specifically, if we let $\mathcal{M}_{\sigma,p}$ be the set of all ergodic σ -invariant measures supported on points fixed by σ^p , we investigate the quantity

$$E_f(p) = \sup_{\nu \in \mathcal{M}_\sigma} \int f d\nu - \sup_{\nu \in \mathcal{M}_{\sigma,p}} \int f d\nu,$$

introduced by G. Yuan and B.R. Hunt [11], being the difference between the maximum integral $\beta(f)$ of f and the ‘best approximating’ periodic orbit whose period divides p .

Our motivation is twofold. Firstly, the above problem is of interest in control theory: if we are able to control the dynamical system $\sigma: \Sigma_A \rightarrow \Sigma_A$, we will wish to direct the trajectories of points in Σ_A toward particular unstable orbits which optimise some given quantity. For reasons of computational simplicity, it is desirable that such orbits should be periodic and have small length. This leads us to ask how close to optimal orbits of a specified period may be. These considerations motivate the work of G. Yuan, B. R. Hunt and E. Ott [7, 11].

Secondly, the quantity $\beta(f)$ has shown itself to be of interest in a number of situations in ergodic theory, both intrinsically [1, 2, 6, 8] and in application to existing problems [3, 4, 9]. It is therefore of interest to be able to compute this quantity accurately in numerical experiments. One obvious approach to this task would be

to exhaustively compute ergodic averages of f along periodic orbits of length up to n , and take the supremum of these averages as an approximation to $\beta(f)$. The error incurred in this approximation would therefore equal $\inf_{1 \leq p \leq n} E_p(f)$.

It is classical [10] that $\cup_{p=1}^{\infty} \mathcal{M}_{\sigma,p}$ is dense in \mathcal{M}_{σ} in the weak-* topology, and so $E_f(p) \rightarrow 0$ as $p \rightarrow \infty$ for all continuous f . We wish to investigate the rate at which this convergence occurs when f is Hölder.

The behaviour of $E_f(p)$ as $p \rightarrow \infty$ is at present poorly understood. On the strength of numerical experiments [7] combined with rigorous analysis, Yuan and Hunt [11] observed that the sequence $E_f(p)$ often decays to zero at an exponential rate, but were unable to prove this in generality. They then asked whether it could be shown that $E_f(p)$ always decays exponentially.

The purpose of this paper is to answer this question in the negative. We have

Theorem 1. *Let $\sigma: \Sigma_A \rightarrow \Sigma_A$ be a subshift of finite type. There exists Hölder continuous $f: \Sigma_A \rightarrow \mathbb{R}$ such that $E_f(p)$ tends to 0 at a slower than stretched-exponential rate: $\log E_f(p) = o(p^\varepsilon)$ for every $\varepsilon > 0$.*

We note that a result similar to Theorem 1 has recently been proved by X. Bressaud and A. Quas in [5], in which the quantity $\inf_{1 \leq p \leq n} E_p(f)$ is considered. Bressaud and Quas are able to obtain upper and lower bounds which are superior to those in the present article, but with the weakness that their lower bound applies only along subsequences of integers n .

2. NOTATION AND DEFINITIONS

Let A be an irreducible aperiodic $N \times N$ matrix of zeroes and ones. We define the *shift space* associated to A to be the set

$$\Sigma_A := \{x = (x_i)_{i \geq 1} : x_i \in \{1, \dots, N\} \text{ and } A(x_i, x_{i+1}) = 1 \text{ for all } i \geq 1\},$$

and define the *shift map* $\sigma: \Sigma_A \rightarrow \Sigma_A$ by

$$(\sigma x)_i = x_{i+1}$$

for all $i \geq 1$. Given $\theta \in (0, 1)$ we define the θ -metric d_θ on Σ_A by

$$d_\theta \left((x_i)_{i \geq 1}, (y_i)_{i \geq 1} \right) = \theta^{\inf\{n \geq 1 : x_n \neq y_n\}}.$$

We say that a function $f: \Sigma_A \rightarrow \mathbb{R}$ is θ -Hölder continuous if it is Lipschitz continuous with respect to the metric d_θ . We fix $\theta \in (0, 1)$ for the remainder of this paper.

We define a *finite word* to be a finite sequence $\omega = (\omega_i)_{i=1}^n$ taking values in the set $\{1, 2, \dots, N\}$. We say that $\omega = (\omega_i)_{i=1}^n$ is *compatible with the matrix A* if $A(\omega_i, \omega_{i+1}) = 1$ for all $i < n$. We define the *length* of the word $(\omega_i)_{i=1}^n$ to be n . We will on occasion describe elements of Σ_A as *infinite words*. We say that finite words ω^1, ω^2 of equal length n are *rotation equivalent* if there exists a nonnegative integer $r < n$ such that $\omega_i^1 = \omega_{i+r}^2$ whenever $1 \leq i \leq n - r$ and $\omega_i^1 = \omega_{i+r-n}^2$ whenever $n - r < i \leq n$. When ω^1 and ω^2 are rotation equivalent we write $\omega^1 \simeq \omega^2$. Words of distinct lengths are never rotation equivalent.

Given two words ω^1, ω^2 with lengths n_1 and n_2 , we define their *concatenation* $\omega^1 \cdot \omega^2$ to be the word of length $n_1 + n_2$ given by $[\omega^1 \cdot \omega^2]_i = \omega_i^1$ when $1 \leq i \leq n_1$, and $[\omega^1 \cdot \omega^2]_i = \omega_i^2$ when $n_1 + 1 \leq i \leq n_1 + n_2$. Given a finite list of words $\omega^1, \dots, \omega^m$, we denote the compound concatenation $\omega^1 \cdot \omega^2 \cdot \dots \cdot \omega^m$ by $\prod_{k=1}^m \omega^k$. Concatenation is

associative. If ω^1, ω^2 are finite words with lengths n_1, n_2 compatible with A , their concatenation is compatible with A if and only if $A(\omega_{n_1}^1, \omega_1^2) = 1$.

For each $p > 0$ we let Ω_p be the set of all words of length p which are compatible with A . We let $\overline{\Omega}_p$ be the set of all words $\omega \in \Omega_p$ such that $A(\omega_p, \omega_1) = 1$. Note that $\omega \in \overline{\Omega}_p$ if and only if both $\omega \in \Omega_p$ and $\omega \cdot \omega \in \Omega_{2p}$.

If $x \in \Sigma_A$ it is clear that $\sigma^p x = x$ if and only if $x_i = x_{i+p}$ for all $i \geq 1$, if and only if there exists $\omega \in \overline{\Omega}_p$ such that $x_{i+kp} = \omega_i$ for all $1 \leq i \leq p$ and $k \geq 0$. In this case we write $x = \pi(\omega)$. This defines a relationship between the sets $\text{Fix}_p = \{x \in \Sigma_A : \sigma^p x = x\}$ and $\overline{\Omega}_p$ which may readily be seen to be bijective. Moreover, we have $\pi(\omega^1) = \sigma^j \pi(\omega^2)$ for some $j \geq 0$ if and only if $\omega^1 \simeq \omega^2$.

If $a = (a_i)_{i=1}^n$ is a finite word and b is either a finite word or an element of Σ_A , we write $a \prec b$ if there is $k \geq 0$ such that $a_i = b_{i+k}$ for all $1 \leq i \leq n$. In this case we say that a is a *subword* of b .

3. PROOF OF THEOREM 1

We begin with the following:

Proposition 3.1. *Let $K = \sigma K$ be a closed nonempty subset of Σ_A and define $f_K(x) = -d_\theta(K, x)$ for all $x \in \Sigma_A$. Clearly f_K is θ -Hölder continuous. For each $\omega \in \Omega_p$ we define*

$$\xi(\omega, K) := \inf_{\omega' \simeq \omega} \{\ell > 0 : \exists a \in \Omega_\ell \text{ such that } a \prec \omega' \text{ and } \forall x \in K, a \not\prec x\}.$$

Let $p > 0$, and suppose that $\sup_{\omega \in \Omega_p} \xi(\omega, K) \leq p$. Then

$$E_{f_K}(p) \geq \frac{1}{p} \theta^{\sup_{\omega \in \Omega_p} \xi(\omega, K)} = \frac{1}{p} \inf_{\omega \in \Omega_p} \theta^{\xi(\omega, K)}.$$

Proof. Let $p > 0$ and $\omega \in \Omega_p$, where $\xi(\omega, K) \leq p$. We have

$$\begin{aligned} \log_\theta \sup_{\omega' \simeq \omega} d(\pi(\omega'), K) &= \log_\theta \sup_{\omega' \simeq \omega} \inf_{x \in K} d_\theta(\pi(\omega'), x) \\ &= \inf_{\omega' \simeq \omega} \sup_{x \in K} \inf \{\ell > 0 : x_\ell \neq \pi(\omega')_\ell\} \\ &= \inf_{\omega' \simeq \omega} \sup \{\ell > 0 : \exists x \in K \text{ s.t. } x_i = \pi(\omega')_i \forall 1 \leq i < \ell\} \\ &= \inf_{\omega' \simeq \omega} \{\ell > 0 : \exists a \in \Omega_\ell \text{ s.t. } a_i = \pi(\omega')_i \forall 1 \leq i \leq \ell \text{ and } \forall x \in K, a \not\prec x\} \\ &= \inf_{\omega' \simeq \omega} \{\ell > 0 : \exists a \in \Omega_\ell \text{ s.t. } a \prec \pi(\omega') \text{ and } a \not\prec x \forall x \in K\} \\ &= \inf_{\omega' \simeq \omega} \{\ell > 0 : \exists a \in \Omega_\ell \text{ s.t. } a \prec \omega' \text{ and } a \not\prec x \forall x \in K\} \\ &= \xi(\omega, K), \end{aligned}$$

where we have used $\xi(\omega, K) \leq p$ in the second-from-last equality. Since K is closed, σ -invariant and nonempty, the Krylov-Bogolioubov Theorem shows that there exists

$\mu \in \mathcal{M}_\sigma$ such that $\mu(K) = 1$. It follows that $\beta(f) = 0$, and so

$$\begin{aligned} E_{f_K}(p) &= \inf_{\sigma^p x = x} \frac{1}{p} \sum_{j=0}^{p-1} d(\sigma^j x, K) \\ &\geq \frac{1}{p} \inf_{\sigma^p x = x} \sup_{0 \leq j < p} d(\sigma^j x, K) \\ &= \frac{1}{p} \inf_{\omega \in \overline{\Omega}_p} \sup_{\omega' \simeq \omega} d(\pi(\omega'), K) \\ &\geq \frac{1}{p} \inf_{\omega \in \Omega_p} \theta^{\xi(\omega, K)}, \end{aligned}$$

as required. \square

It follows that to prove Theorem 1, it suffices to construct a nonempty compact set $K = \sigma K \subseteq \Sigma_A$ such that $\sup_{\omega \in \Omega_p} \xi(\omega, K) = o(p^\varepsilon)$ as $p \rightarrow \infty$. The remainder of this section is dedicated to this task. We will construct the invariant set K recursively, using a sequence of sets of words M_n of increasing length. The set K will then arise as a limit of these sets.

Let M_1 be a subset of Ω_{ℓ_1} for some positive integer ℓ_1 such that $m_1 = \text{Card } M_1$ is divisible by 2^{16} . We require that M_1 have the following properties:

Definition 3.2. *There exists an integer $P \geq 1$ and word $z = (z_i)_{i=1}^P \in \Omega_P$ such that:*

- (i) *Every $\omega \in M_1$ satisfies $z_i = \omega_i$ for all $1 \leq i \leq P$.*
- (ii) *If $\omega = \omega^1 \cdot \omega^2$ where $\omega^1, \omega^2 \in M_1$, and $z_i = \omega_{i+r}$ for all $1 \leq i \leq P$, then either $r = \ell_1$ or $r = 0$.*
- (iii) *$A(\omega_1^1, \omega_1^2) = 1$ for all $\omega^1, \omega^2 \in M_1$; that is, $\omega^1 \cdot \omega^2 \in \Omega_{2\ell_1}$ for every pair $\omega^1, \omega^2 \in M_1$.*

The reader may verify that such a set M_1 may be constructed for any prescribed m_1 and matrix A .

We will define a sequence of sets M_n with cardinalities m_n consisting of words of length ℓ_n in an inductive fashion starting with the set M_1 . We begin by defining some integer sequences which will be crucial to our construction.

Definition 3.3. *Given an integer m_1 divisible by 2^{16} , let $q_1 = t_1 = m_1/4$. Define sequences $(m_n)_{n \geq 1}$, $(q_n)_{n \geq 1}$ and $(t_n)_{n \geq 1}$ as follows. Given the integer t_n , we let $m_{n+1} = 4^{t_n}$ and $q_{n+1} = \frac{1}{4}m_{n+1} = 4^{t_n-1}$, and let τ_{n+1} be the unique positive real number such that*

$$\tau_{n+1}^{\sqrt{\tau_{n+1}}} = 4^{t_n-1} = q_{n+1}.$$

We then define

$$t_{n+1} = 4 \left\lfloor \frac{t_n}{\sqrt{\tau_{n+1}}} \right\rfloor^{-1}.$$

The key properties of these sequences are stated in the lemma below.

Lemma 3.4. *For each $n \geq 1$, we have $t_n \mid q_n$, $m_n/q_n = 4$ and $t_n \geq 2^{14}$. Moreover, the sequence t_n satisfies $t_n/\sqrt{t_{n+1}} \leq 16t_n^{1/4}$ for all $n \geq 1$ and*

$$\lim_{n \rightarrow \infty} \frac{t_n}{t_{n+1}} = 0.$$

Proof. The first two statements are clear. We consider the sequence $(t_n)_{n \geq 1}$; the definition implies that

$$\frac{1}{16} \tau_{n+1} = 4 \frac{t_n - 1}{\sqrt{\tau_{n+1}}} - 2 \leq 4 \left\lfloor \frac{t_n}{\sqrt{\tau_{n+1}}} \right\rfloor - 1 = t_{n+1} \leq 4 \frac{t_n}{\sqrt{\tau_{n+1}}} - 1 \leq \tau_{n+1},$$

for every $n \geq 1$. We proceed inductively. Given that $t_n \geq 2^{14}$, we notice that

$$\sqrt{\tau_{n+1}} \log \tau_{n+1} = (t_n - 1) \log 4.$$

We thus have

$$t_n = \sqrt{\tau_{n+1}} \log_4 \tau_{n+1} + 1 > \sqrt{\tau_{n+1}},$$

and therefore

$$\log_4 \tau_{n+1} < 2 \log_4 t_n.$$

We deduce

$$\sqrt{t_{n+1}} \geq \frac{1}{4} \sqrt{\tau_{n+1}} = \frac{t_n - 1}{4 \log_4 \tau_{n+1}} \geq \frac{t_n - 1}{8 \log_4 t_n} \geq \frac{1}{8} t_n^{3/4} - \frac{1}{8},$$

where we have used the elementary inequality $\log_4 t \leq t^{1/4}$ for all $t \geq 16$. Thus

$$t_{n+1} \geq \frac{1}{64} \left(t_n^{3/4} - 1 \right)^2 \geq 2^{-6} \left(2^{21} - 2^{11} \sqrt{2} + 1 \right) > 2^{14}$$

for every $n \geq 1$, making it clear that $t_n / \sqrt{t_{n+1}} \leq 16 t_n^{1/4}$. One may easily use the above to show that $t_{n+1} \geq t_n + \frac{1}{64}$ for each $n \geq 2$, implying that $t_n \rightarrow \infty$; since for each n we have

$$0 \leq \frac{t_n}{t_{n+1}} \leq \frac{4t_n}{t_n^{3/2} - 2t_n^{3/4} + 1},$$

it follows that $\lim_{n \rightarrow \infty} t_n / t_{n+1} = 0$ as required. \square

For an integer $n \geq 1$, a finite word $a = (a_i)_{i=1}^m$ and a finite or infinite word b , we shall write $a \prec_n b$ if there is $k \geq 0$ such that $a_i = b_{i+k\ell_n}$ for all $1 \leq i \leq m$. The distinction between \prec and \prec_n will be important since we will construct words $\omega \in M_{n+1}$ as concatenations of words $a, b, c, \dots \in M_n$. For example, if $a, b, c \in M_n$, then it is true that $b \cdot c \prec_n a \cdot b \cdot c \cdot a$ and $c \cdot b \not\prec_n a \cdot b \cdot c \cdot a$; however, the statement $c \cdot b \prec a \cdot b \cdot c \cdot a$ could be either true or false, depending on the subword structure of the words a, b and c .

Given $n \geq 1$ and the set M_n , we construct the set M_{n+1} as follows. Recalling that $m_n = 4q_n$, partition M_n into q_n disjoint sets C_n^k of cardinality 4, where $1 \leq k \leq q_n$. For each k , we will write $C_n^k = \{c_j^k : 1 \leq j \leq 4\}$. Define

$$I_n = \{(i_1, i_2, \dots, i_{q_n}) \in \{1, 2, 3, 4\}^{q_n} : i_m = i_{m+t_n} \text{ for all } 1 \leq m \leq q_n - t_n\}.$$

For each q_n -tuple $(i_1, \dots, i_{q_n}) \in I_n$, we construct the word

$$\omega_{(i_1, \dots, i_{q_n})} := \prod_{k=1}^{q_n} c_{i_k}^k.$$

We let $M_{n+1} = \{\omega_{(i_1, \dots, i_{q_n})} : (i_1, \dots, i_{q_n}) \in I_n\}$. Clearly,

$$\text{Card } M_{n+1} = \text{Card } I_n = 4^{t_n} = m_{n+1}$$

in accordance with Definition 3.4. We remark that Definition 3.2(iii) implies that $M_n \subseteq \Omega_{\ell_n}$ for every $n \geq 1$. The key features of the above construction are summarised in the following lemma.

Lemma 3.5. *The following are direct consequences of the definition of the sets M_n :*

- (i) *If $\prod_{k=1}^{q_n} c_{j_k}^k \in M_n$, where $n \geq 2$ and each $c_{j_k}^k \in C_{n-1}^k$, then $c_{j_k}^k \in C_{n-1}^1$ if and only if $k = 1$.*
- (ii) *If $\omega_1 = c_{j_1}^{k_1} \cdot c_{j_2}^{k_2}$ with $c_{j_1}^{k_1} \in C_n^{k_1}, c_{j_2}^{k_2} \in C_n^{k_2}$, then there exists $N > n$ and $\omega_2 \in M_N$ such that $\omega_1 \prec_n \omega_2$ if and only if $k_2 = 1 + k_1 \pmod{q_n}$.*
- (iii) *If $\omega_1 = \prod_{k=k_0}^{k_0+t_n} c_{j_k}^k$ where each $c_{j_k}^k \in C_n^k$, and $k_0 \leq q_n - t_n$, and $j_{k_0} \neq j_{k_0+t_n}$, then $\omega_1 \not\prec_n \omega_2$ for all $\omega_2 \in M_N$, for all $N \geq n + 1$.*
- (iv) *If $\omega_1 \prec_n \omega_2 \in M_{n+1}$, $\omega_1 \prec_n \omega_3 \in M_{n+1}$ and ω_1 has length at least $t_n \ell_n$, then $\omega_2 = \omega_3$.*
- (v) *For each $n \geq 1$ we have*

$$\ell_{n+1} = q_n \ell_n = \frac{1}{4} m_n \ell_n$$

and hence

$$\ell_{n+1} = \ell_1 \prod_{k=1}^n q_k.$$

The proof is clear. \square

The following lemma allows us to pass from the relation \prec to the relation \prec_n , and thus make use of Lemma 3.5.

Lemma 3.6. *Let $a = (a_i)_{i=1}^{\ell_n} \in M_n$ and $\omega = (\omega_i)_{i=1}^{\ell_N} \in M_N$ where $n < N$. Suppose that there is $r \geq 0$ such that $a_i = \omega_{r+i}$ for all $1 \leq i \leq \ell_n$. Then $\ell_n \mid r$.*

Proof. We first prove the case $n = 1$. Let $\omega = \prod_{k=1}^m \omega^k$ where each $\omega^k \in M_1$, and suppose $a_i = \omega_{r+i}$ for all $1 \leq i \leq \ell_1$. There exists k^* such that $k^* \ell_1 \leq r + i < (k^* + 2) \ell_1$ for all $1 \leq i \leq \ell_1$, so that if we let $\hat{\omega} = \omega^{k^*} \cdot \omega^{k^*+1}$ then $z_i = a_i = \hat{\omega}_{i+r-k^* \ell_1}$ for all $1 \leq i \leq P \leq \ell_1$ by Definition 3.2 (i). By Definition 3.2 (ii) we have either $r - k^* \ell_1 = \ell_1$ or $r - k^* \ell_1 = 0$ and so $\ell_1 \mid r$ as required.

We proceed by induction on n . Let $a = \prod_{k=1}^{q_n-1} \hat{a}^k$ and $\omega = \prod_{k=1}^m \hat{\omega}^k$ with each $\hat{a}^k, \hat{\omega}^k \in M_{n-1}$. Let $\hat{a}^1 = (\hat{a}^1)_{i=1}^{\ell_{n-1}}$. Since $\hat{a}_i^1 = \omega_{r+i}$ for all $1 \leq i \leq \ell_{n-1}$, we have $\ell_{n-1} \mid r$ by the induction hypothesis. Since each $\hat{\omega}^k$ has length ℓ_{n-1} , it follows that there is $s = r/\ell_{n-1} > 0$ such that $\prod_{k=1}^{q_n-1} \hat{a}^k = \prod_{k=s}^{k=s+q_n-1} \hat{\omega}^k$. By Lemma 3.5(i) we have $\hat{a}^k \in C_{n-1}^1$ if and only if $k = 1$, and similarly $\hat{\omega}^k \in C_{n-1}^1$ if and only if $k \equiv 1 \pmod{q_n}$. Since $\hat{\omega}^{s+1} \in C_{n-1}^1$ it follows that $s \equiv 0 \pmod{q_n}$. Since $r = s \ell_{n-1}$ and $\ell_n = q_{n-1} \ell_{n-1}$ we deduce that $\ell_n \mid r$. \square

Henceforth, we shall say that a finite word ω is *admissible* if and only if it occurs as a subword of some $\omega_n \in M_n$ for some n (hence for all sufficiently large n). We deduce:

Corollary 3.7. *Let $\omega = (\omega_i)_{i=1}^p$ be admissible. Let $a, b \in M_n$, and write $a = (a_i)_{i=1}^{\ell_n}, b = (b_i)_{i=1}^{\ell_n}$. If there are $r, s > 0$ such that $a_i = \omega_{i+r}, b_i = \omega_{i+s}$ for all $1 \leq i \leq \ell_n$, then $\ell_n \mid r - s$.*

We may now prove:

Lemma 3.8. *The following constraints on admissibility hold:*

- (i) *If $\omega = c_{j_1}^{k_1} \cdot c_{j_2}^{k_2}$ with $c_{j_1}^{k_1} \in C_n^{k_1}, c_{j_2}^{k_2} \in C_n^{k_2}$, then ω is admissible if and only if $k_2 = 1 + k_1 \pmod{q_n}$.*

(ii) If $\omega = \prod_{k=k_0}^{k_0+t_n} c_{j_k}^k$ with each $c_{j_k}^k \in C_n^k$, and $j_{k_0} \neq j_{k_0+t_n}$, and $k_0 \leq q_n - t_n$, then ω is not admissible.

Proof. Apply Lemma 3.5 and Lemma 3.6. \square

Lemma 3.9. *Let $a, b \in M_n$ where $n \geq 2$, and suppose that the word $a \cdot b$ is inadmissible. Then $a \cdot b$ has an inadmissible subword of length less than or equal to $2t_{n-1}\ell_{n-1}$.*

Proof. Let $a = \prod_{k=1}^{q_{n-1}} a^k$, $b = \prod_{k=1}^{q_{n-1}} b^k$ where each $a^k, b^k \in M_{n-1}$. Define $u = \prod_{k=q_{n-1}-t_{n-1}+1}^{q_{n-1}} a^k$ and $v = \prod_{k=1}^{t_{n-1}} b^k$, each being an admissible word of length $t_{n-1}\ell_{n-1}$. Clearly $u \cdot v \prec a \cdot b$. Suppose, for a contradiction, that $u \cdot v \prec \omega_1 \cdot \omega_2 = \omega$, say, where $\omega_1, \omega_2 \in M_n$ and ω is admissible. Let $(u \cdot v)_i = \omega_{i+r}$ for all $1 \leq i \leq 2\ell_{n-1}$. By Lemma 3.6, $r = \ell_{n-1}s$ for some s . Suppose that $s \geq q_n$; then $v \prec_n \omega_2$ and so $\omega_2 = b$ by Lemma 3.5(iv). Thus $s = q_n$ and so $u \prec_n \omega_1$ and $\omega_1 = a$. Thus $\omega = a \cdot b$ and so is not admissible. It follows that $u \cdot v$ is inadmissible; this word has length $2t_{n-1}\ell_{n-1}$ as required. \square

Lemma 3.10. *Let ω be a word of length $\ell \geq \ell_n$, where $n \geq 2$. Then either $\omega \simeq \prod_{k=1}^m \omega^k$ with each $\omega^k \in C_{n-1}^k$, or there exists $\omega' \simeq \omega$ which has an inadmissible subword of length less than or equal to $3\ell_{n-1}$.*

Proof. Suppose that for all $\omega' \simeq \omega$, every subword of ω' of length $3\ell_{n-1}$ is admissible. Let $\ell = s\ell_{n-1} + r$ with $0 \leq r < \ell_{n-1}$. We claim that there exist $\omega^k \in M_{n-1}$ and a word ω^* of length r such that $\omega \simeq \prod_{k=1}^s \omega^k \cdot \omega^*$. We will show that $(\omega_i)_{i=1}^{m\ell_{n-1}} \simeq \prod_{k=1}^m \omega^k$ for all $m \leq s$ by induction on m .

Clearly, any admissible word of length $3\ell_{n-1}$ must include some $a \in M_{n-1}$ as a subword, and so ω must include such a subword. Taking a rotation equivalent of ω if necessary, we deduce that there exists $\omega^1 \in M_{n-1}$ such that $\omega_i = \omega_i^1$ for all $1 \leq i \leq \ell_{n-1}$. This proves the case $m = 1$.

Given that $(\omega_i)_{i=1}^{m\ell_{n-1}} = \prod_{k=1}^m \omega^k$ with each $\omega^k \in M_{n-1}$, consider the word $b = (b_i)_{i=1}^{2\ell_{n-1}}$ defined by $b_i = \omega_{(m-1)\ell_{n-1}+i}$ for all $1 \leq i \leq 2\ell_{n-1}$, which is well-defined as long as $m+1 \leq s$. Since b has length $2\ell_{n-1} < 3\ell_{n-1}$ it is admissible, and so there exist $N, M, t > 0$ and $a = \prod_{k=1}^M a^k \in M_N$ with each $a^k \in M_{n-1}$ such that $b_i = a_{t+i}$ for all $1 \leq i \leq 2\ell_{n-1}$. By Lemma 3.6 we have $t = \ell_{n-1}\bar{t}$ for some \bar{t} , so that $b = a^{\bar{t}+1} \cdot a^{\bar{t}+2}$. Thus $(\omega_i)_{i=m\ell_{n-1}+1}^{(m+1)\ell_{n-1}} = (b_i)_{i=\ell_{n-1}+1}^{2\ell_{n-1}} = a^{\bar{t}} \in M_{n-1}$ as required to prove the case $m+1$. This completes the induction step and proves our claim.

Since for every $\omega' \simeq \omega$, every subword of ω' with length less than or equal to $3\ell_{n-1}$ is admissible, the word $z = \omega^s \cdot \omega^* \cdot \omega^1$ must be admissible. By Corollary 3.7 this implies $\ell_{n-1} \mid r$ and hence $r = 0$. This completes the proof. \square

Lemma 3.11. *Let $\omega = \prod_{k=1}^m \omega^k$ for some $\omega^1, \dots, \omega^m \in M_n$ with $m \geq 1$. Then at least one of the following holds:*

- (a) *There exists $\omega' \simeq \omega$ such that ω' has an inadmissible subword with length less than or equal to $2\ell_{n-1}t_{n-1}$, or*
- (b) *there exists $\omega' \simeq \omega$ such that ω' has an inadmissible subword with length less than or equal to $(1+t_n)\ell_n$, and $m \geq q_n$, or*
- (c) *there exist $\hat{m} > 0$ and $\hat{\omega}^1, \dots, \hat{\omega}^{\hat{m}} \in M_{n+1}$ such that $m = \ell_{n+1}\hat{m}$ and $\omega \simeq \prod_{k=1}^{\hat{m}} \hat{\omega}^k$.*

Proof. Suppose that the case (a) does not hold. Writing $\omega = \prod_{i=1}^m c_{j_i}^{k_i}$ with every $c_{j_i}^{k_i} \in C_n^{k_i}$, this assumption implies via Lemma 3.8(i) and Lemma 3.9 that for every $i < m$ we have $k_{i+1} = 1 + k_i \pmod{q_n}$. Clearly, there exists a rotation equivalent of ω which includes the word $c_{j_m}^{k_m} \cdot c_{j_1}^{k_1}$. Again by Lemma 3.8(i) and Lemma 3.9, our assumption that case (a) does not hold implies that this word is admissible, and hence $k_1 = k_m + 1 \pmod{q_n}$. It follows that the sequence k_i must take every value in the range $1, \dots, q_n$ an equal number of times. Taking a rotation equivalent if necessary, we have

$$\omega = \prod_{s=1}^S \prod_{k=1}^{q_n} c_{j_{k,s}}^k,$$

for some integer $S \geq 1$ and some sequence $j_{k,s}$. Note in particular that $m \geq q_n$ (as required for (b) to hold).

We now suppose also that (b) does not hold. To show that ω is a concatenation of elements of M_{n+1} , and hence that (c) holds, it remains to show that $j_{k,s} = j_{k+t_n,s}$ for all $1 \leq s \leq S$ and $1 \leq k \leq q_n - t_n$. If this is not the case, then ω must include a subword of the form

$$\prod_{k=k_0}^{k_0+t_n} c_{j_{k,s}}^k$$

for some fixed s such that $j_{k_0,s} \neq j_{k_0+t_n,s}$, which is inadmissible by Lemma 3.8(ii) and has length $(1+t_n)\ell_n$, implying the case (b). \square

Combining the above lemmata, we obtain:

Proposition 3.12. *Define a set $K \subseteq \Sigma_A$ by letting $x \in K$ if and only if every subword of x is admissible. Then K is closed, satisfies $\sigma K = K$, and is nonempty. Moreover,*

$$\sup_{\omega \in \Omega_p} \xi(\omega, K) = o(p^\varepsilon),$$

for every $\varepsilon > 0$.

Proof. That K is a nonempty subset of Σ_A follows from the fact that $M_n \subseteq \Omega_{\ell_n}$ for every $n \geq 1$; the proof that K is closed and $\sigma K = K$ is straightforward. Given $\omega \in \Omega_p$ and $\ell_n \leq p < \ell_{n+1}$, we will attempt to bound the quantity

$$\xi(\omega, K) = \inf_{\omega' \simeq \omega} \{ \ell > 0 : \exists a \in \Omega_\ell \text{ such that } a \prec \omega' \text{ and } a \text{ is not admissible} \}.$$

Suppose that $\xi(\omega, K) > 3\ell_{n-1}$. Then by Lemma 3.10 we have $\omega \simeq \prod_{i=1}^m c_{j_i}^{k_i}$ where each $c_{j_i}^{k_i} \in C_{n-1}^{k_i}$ and $m = p/\ell_{n-1}$. Lemma 3.11 then implies that either $\xi(\omega, K) \leq 2\ell_{n-2}t_{n-2}$, or $\xi(\omega, K) \leq \ell_{n-1}(1+t_{n-1})$, or $\omega \simeq \prod_{i=1}^{\hat{m}} c_{j_i}^{k_i}$ where each $c_{j_i}^{k_i} \in C_n^{k_i}$ and $\hat{m} = p/\ell_n$. In the last of these three cases, we apply Lemma 3.11 again to see that, since $p < \ell_{n+1}$, either $\xi(\omega, K) \leq 2\ell_{n-1}t_{n-1}$, or $p \geq \ell_n q_n$ and $\xi(\omega, K) \leq \ell_n(1+t_n) \leq 2pt_n/q_n$. We conclude that in any case

$$\begin{aligned} \xi(\omega, K) &\leq \max \left\{ 3\ell_{n-1}, 2\ell_{n-2}t_{n-2}, \ell_{n-1}(t_{n-1} + 1), 2\ell_{n-1}t_{n-1}, \frac{2pt_n}{q_n} \right\} \\ &= \max \left\{ 2\ell_{n-1}t_{n-1}, \frac{2pt_n}{q_n} \right\}. \end{aligned}$$

Hence for every $n \geq 3$ and $\varepsilon > 0$,

$$\begin{aligned} \sup_{\ell_n \leq p < \ell_{n+1}} \sup_{\omega \in \Omega_p} \frac{1}{p^\varepsilon} \xi(\omega, K) &\leq \max \left\{ 2\ell_{n-1}t_{n-1}\ell_n^{-\varepsilon}, 2\frac{\ell_{n+1}^{1-\varepsilon}t_n}{q_n} \right\} \\ &\leq 2 \max \{ \ell_{n-1}^{1-\varepsilon}t_{n-1}q_{n-1}^{-\varepsilon}, \ell_n^{1-\varepsilon}t_nq_n^{-\varepsilon} \} = 2\ell_n^{1-\varepsilon}t_nq_n^{-\varepsilon}. \end{aligned}$$

To complete the proof, therefore, it suffices to show that $t_nq_n^{-\varepsilon}\ell_n \rightarrow 0$ for every $\varepsilon > 0$. Using Lemma 3.5(vii), Lemma 3.4 and Definition 3.3, we have

$$\begin{aligned} \log_4(t_nq_n^{-\varepsilon}\ell_n) &\leq \log_4 \left(4^{\frac{t_{n-1}}{\sqrt{t_n}} - \varepsilon t_{n-1}} q_1 \ell_1 \prod_{k=2}^{n-1} q_k \right) \\ &\leq \log_4(q_1 \ell_1) + \frac{t_{n-1}}{\sqrt{t_n}} - \varepsilon t_{n-1} + \sum_{k=1}^{n-2} t_k \\ &= -\varepsilon t_{n-1} + \sum_{k=1}^{n-2} t_k + O\left(t_{n-1}^{1/4}\right). \end{aligned}$$

Elementary analysis then shows that since t_{n-1}/t_n tends to zero, $\sum_{k=2}^{n-2} t_k = o(t_{n-1})$ and so

$$\log_4(t_nq_n^{-\varepsilon}\ell_n) \rightarrow -\infty$$

as required. □

REFERENCES

- [1] J. BRÉMONT, ‘Gibbs measures at temperature zero’, *Nonlinearity* 16 (2003), 419-426.
- [2] T. BOUSCH, ‘Le poisson n’a pas d’arêtes’, *Ann. Inst. H. Poincaré Probab. Statist.* 36 (2000) 489-508.
- [3] T. BOUSCH and O. JENKINSON, ‘Cohomology classes of dynamically non-negative C^k functions’, *Invent. Math.* 148 (2002) 207-217.
- [4] T. BOUSCH and J. MAIRESSE, ‘Asymptotic height optimization for topological IFS, Tetris heaps, and the finiteness conjecture’, *J. Amer. Math. Soc.*, 15 (2002) 77-111.
- [5] X. BRESSAUD and A. QUAS, ‘Rates of approximation of minimizing measures’, *Nonlinearity* 20 (2007) 845-853.
- [6] G. CONTRERAS, A. LOPES and P. THIEULLEN, ‘Lyapunov minimizing measures for expanding maps of the circle’, *Ergod. Theory Dynam. Systems* 5 (2001) 1379-1409.
- [7] B. HUNT and E. OTT, ‘Optimal periodic orbits of chaotic systems occur at low period’, *Phys. Rev. E* 54 (1996) 328-337.
- [8] O. JENKINSON, ‘Ergodic optimization’, *Disc. Cont. Dyn. Syst.* 15 (2006) 197-224.
- [9] O. JENKINSON, ‘Rotation, entropy and equilibrium states’, *Trans. Amer. Math. Soc* 353 (2001), 3713-3739.
- [10] K.R. PARTHASARATHY, ‘On the category of ergodic measures’, *Illinois J. Math.* 5 (1961) 648-656.
- [11] G. YUAN and B. HUNT, ‘Optimal orbits of hyperbolic systems’, *Nonlinearity* (4) 12 (1999) 1207-1224.

D. COLLIER, SCHOOL OF MATHEMATICS, UNIVERSITY OF MANCHESTER, MANCHESTER M13 9PL, U.K.

I. D. MORRIS, MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, U.K.